

Liver Tonic

NATURAL HOMEOPATHICS



Item# VHT-LT
4 Fl. Oz

Product Indication: Provides homeopathic support and drainage for liver function.

Ingredients: HPUS; Phosphorus 6X, Chelidonium Majus 2X, Liver 3X, 6X, Lycopodium Clavatum 4X, Hydrastis Canadensis 2X, Carduus Marianus 2X, Raphanus Sativus 3X, Taraxacum Officinale 2X, Ptelea Trifoliata 2X, Gallbladder 3X, 6X, Byronia Alba 6X, Berberis Vulgaris 4X.

Phosphorus 6X

- Causes yellow atrophy of the liver and sub-acute hepatitis
- Disorganizes the blood, causing fatty degeneration of blood vessels and every tissue and organ of the body
- Gives rise to hemorrhages and jaundice

Chelidonium Majus 2X

- Jaundice skin
- Constant pain under the inferior angle of right scapula
- Enlarged liver
- Profuse, foaming urine

Liver 3X, 6X

- Sarcodes for tissue support

Lycopodium Clavatum 4X

- Failures of digestive powers, where the function of the liver is seriously disturbed.
- Ascities, (an accumulation of serous fluid in the peritoneal cavity) in liver disease

Hydrastis Canadensis 2X, 3X

- Liver torpid, tender
- Jaundice

Carduus Marianus 2X

- Action centered in the liver and portal system, causing soreness, pain and jaundice.
- Hemorrhages, especially connected with hepatic disease
- Hyperemia of liver with jaundice
- Cirrhosis with dropsy

Raphanus Sativus 3X

- Produces pain and stitches in liver and spleen
- Increase of bile and salivary secretion

Taraxacum Officiniale 2X

- Liver enlarged and indurated
- Sharp stitches in left side

Ptelea Trifoliata 2X

- Liver sore, swollen, sensitive to pressure
- Aching and heaviness in the region of the liver is greatly aggravated by lying on the left side

Gallbladder 3X, 6X

- Sarcodes for tissue support

Bryonia Alba 6X

- Liver region swollen, sore, tense
- Tenderness of abdominal walls

Berberis Vulgaris

- Stitching pains in front of kidneys extending to liver, spleen, stomach, and groins

Dosage: For adults and children 12 or more years of age: 1 to 2 teaspoons daily, taken with meals or in juice.
Children under 12: half the adult dosage. Infants: Consult physician prior to use.